

Somatic symptoms (sweating, nausea, diarrhea)									
Exaggerated startle response									
Conditions associated with stress (irritable bowel syndrome, headaches)									
Worry may interfere with attention to tasks									
Perfectionism or worry about quality of performance									
Alcohol or substance abuse									
Potential for psychotic features									
Delusions		*							*
Risk for suicidal behavior									
Occupational problems						*			
Academic problems						*			
Difficulty concentrating or sustaining attention to tasks		*				*			
Agitation/restlessness		*				*		*	
Irritability/temper outbursts									
Marital/family problems									
Distractibility		*							
Talking excessively									
Difficulty awaiting turn		*							
Impulsiveness		*							
Poor judgment								*	
Euphoria									
Inflated self-esteem or grandiosity								*	
Decreased need for sleep								*	
Flight of ideas or racing thoughts								*	
Increase in goal-directed activity/excessive planning								*	
Increased sociability									
Excessive spending/foolish investments								*	
Sexual indiscretion/excessive sexual interest								*	
Disorganized or bizarre social behavior									
Disregard for ethical concerns									
Hostility or threats to others									
Labile mood									
Disorganized speech									*
Grossly disorganized or catatonic behavior									*
Potential for assaultive behavior									*
Hallucinations									*
Negative symptoms (affective flattening, avolition)									*

*Unipolar depression or bipolar disorder. †Patients with bipolar disorder with mixed manic episodes present with depressive and hypomanic symptoms.

Reference:

1. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders: Text Revision*. 4th ed. Washington, DC: American Psychiatric Association; 2000.